

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2001 (27.12.2001)

PCT

(10) International Publication Number
WO 01/97809 A2

(51) International Patent Classification: **A61K 31/436**,
45/06, A61P 9/00, 9/10, 25/28

(74) Agents: **MILOWSKY, Arnold, S.**; American Home Products Corporation, Patent Law Dept., Five Giralda Farms, Madison, NJ 07940 et al. (US).

(21) International Application Number: **PCT/US01/19179**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(22) International Filing Date: **14 June 2001 (14.06.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/212,117 **16 June 2000 (16.06.2000)** **US**

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW); Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM); European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR); OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


(71) Applicant: **AMERICAN HOME PRODUCTS CORPORATION** [US/US], Five Giralda Farms, Madison, NJ 07940-0874 (US).

Published:

— *without international search report and to be republished upon receipt of that report*

(72) Inventors: **AZROLAN, Neal, Ivan**; 286 Glenn Avenue, Lawrenceville, NJ 08648 (US). **SEHGAL, Surendra, Nath**; 20711 76th Avenue S.E., Snohomish, WA 98296 (US). **ADELMAN, Steven, Jay**; 342 Windy Run Drive, Doylestown, PA 18901 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/97809 A2

(54) Title: **METHOD OF TREATING CARDIOVASCULAR DISEASE**

(57) Abstract: The invention provides a method of treating or inhibiting cardiovascular, cerebral vascular, or peripheral vascular disease in a mammal in need thereof, which comprises providing said mammal an effective amount of a rapamycin.

METHOD OF TREATING CARDIOVASCULAR DISEASE

BACKGROUND OF THE INVENTION

5 This invention relates the use of a rapamycin in the treatment and inhibition of cardiovascular disease, cerebral vascular disease, and peripheral vascular disease.

Coronary artery disease, the primary form of cardiovascular disease (CVD), is the major cause of death in the United States today, responsible for over 550,000 deaths per year. Cerebrovascular disease is the third leading cause of death in the United States. The etiology of both coronary artery and cerebrovascular diseases is attributed to atherosclerosis. Through its clinical manifestations, atherosclerosis is the major cause of the more than one million heart attacks and approximately 400,000 strokes that occur each year. In addition to the high morbidity and mortality associated with atherosclerosis, it has been estimated that atherosclerosis has cost the United States' economy over \$80 billion each year in lost wages, lost productivity, and medical care costs [Levy, R., Am. Heart J. 110: 1116 (1985)]. A substantial body of evidence has established a relationship between hypercholesterolemia and premature atherosclerosis; the higher the levels of plasma cholesterol, the greater the risk of subsequent heart attack. [Steinberg, D., JAMA 264: 3047 (1991); Lipid Research Clinics Program, JAMA 251: 351 (1984); Rifkind, B., Am. J. Cardiol. 54: 30C (1984)]. However, recent information demonstrates that the atherosclerotic process is far more complicated than a simple correlation with plasma lipid levels, and that there are both systemic and local factors within the vascular wall that play a major role in the progression of this disease [Sulistiyan, Adelman, S.J., Chandrasekaran, A., Jayo, J. and St. Clair, R.W.. Arteriosclerosis and Thrombosis 15: 837, (1995)].

Atherosclerosis is a complex disease that is associated with a variety of etiologic factors. Studies have shown that, of the major factors involved, diet-induced hyperlipidemia and genetic defects or abnormalities in lipoprotein metabolism have received the most attention. The local disease process of atherosclerosis is characterized by the accumulation of lipids in the walls of blood vessels.

- 2 -

Concomitant with lipid accumulation, there is vascular cell damage resulting in dysfunction of the endothelium, smooth muscle proliferation, and matrix deposition. These changes ultimately result in the formation of what is termed "plaque". As these plaques expand and mature, ruptures in their surface can occur, leading to major thrombotic events. This process, which can occur in essentially all of the blood vessels of the body, results in many of the major disease categories of our time, including coronary artery disease, peripheral vascular disease, myocardial infarction and stroke.

10 Recently, it has been discovered that cells of the immune system play a major role in all of the processes of atherosclerosis, and thus the process has been described as a chronic inflammatory-fibroproliferative disease of the vascular wall. The attachment of monocytes and T-lymphocytes to the injured endothelium followed by their migration into the intima is one of the first and most crucial steps in lesion development. The co-localization of CD4+ T-cells and macrophages in the lesion, the abundant expression of HLA Class II molecules and the co-stimulatory molecule CD40 and its ligand (CD40L) indicate a contribution of cell-mediated immunity to atherogenesis. A wide variety of studies in animal models suggest that T- and B-cells, and monocytes and macrophages promote lesion progression, and in fact, are in essential for the development of atherosclerotic lesions. Importantly, the local vascular wall immune contribution continues throughout, participating in both plaque expansion as well as rupture. In addition to the local process in the vessel wall, systemic signs of an inflammatory reaction are also associated with lesion development. Thus plasma levels of C-reactive protein and fibrinogen and the white blood cell count are positively correlated to the risk of cardiovascular disease.

30 Rapamycin is a macrocyclic triene antibiotic produced by Streptomyces hygroscopicus, which was found to have antifungal activity, particularly against Candida albicans, both in vitro and in vivo [C. Vezina et al., J. Antibiot. 28, 721 (1975); S.N. Sehgal et al., J. Antibiot. 28, 727 (1975); H. A. Baker et al., J. Antibiot. 31, 539 (1978); U.S. Patent 3,929,992; and U.S. Patent 3,993,749]. Additionally, rapamycin alone (U.S. Patent 4,885,171) or in combination with picibanil (U.S. Patent 4,401,653) has been shown to have antitumor activity.

- 3 -


The immunosuppressive effects of rapamycin have been disclosed in FASEB 3, 3411 (1989). Cyclosporin A and FK-506, other macrocyclic molecules, also have been shown to be effective as immunosuppressive agents, and are useful in preventing transplant rejection [FASEB 3, 3411 (1989); FASEB 3, 5256 (1989); R. Y. Calne et al., Lancet 1183 (1978); and U.S. Patent 5,100,899]. R. Martel et al. [Can. J. Physiol. Pharmacol. 55, 48 (1977)] disclosed that rapamycin is effective in the experimental allergic encephalomyelitis model, a model for multiple sclerosis; in the adjuvant arthritis model, a model for rheumatoid arthritis; and effectively inhibited the formation of IgE-like antibodies.

Rapamycin is also useful in preventing or treating systemic lupus erythematosus [U.S. Patent 5,078,999], pulmonary inflammation [U.S. Patent 5,080,899], insulin dependent diabetes mellitus [U.S. Patent 5,321,009], skin disorders, such as psoriasis [U.S. Patent 5,286,730], bowel disorders [U.S. Patent 5,286,731], smooth muscle cell proliferation and intimal thickening following vascular injury [U.S. Patents 5,288,711 and 5,516,781], adult T-cell leukemia/lymphoma [European Patent Application 525,960 A1], ocular inflammation [U.S. Patent 5,387,589], malignant carcinomas [U.S. Patent 5,206,018], cardiac inflammatory disease [U.S. Patent 5,496,832], and anemia [U.S. Patent 5,561,138].

DESCRIPTION OF THE INVENTION

This invention provides a method of treating or inhibiting cardiovascular disease or peripheral vascular disease in a mammal in need thereof, which comprises providing an effective amount of a rapamycin to said mammal. As defined herein, the term "a rapamycin" defines a class of immunosuppressive compounds which contain the basic rapamycin nucleus (shown below). The rapamycins of this invention include compounds which may be chemically or biologically modified as derivatives of the rapamycin nucleus, while still retaining immunosuppressive properties. Accordingly, the term "a rapamycin" includes esters, ethers, oximes, hydrazones, and hydroxylamines of rapamycin, as well as rapamycins in which functional groups on the rapamycin nucleus have been modified, for example through reduction or oxidation. The term "a rapamycin" also includes pharmaceutically acceptable salts of rapamycins, which are capable of forming such salts, either by virtue of containing an acidic or basic moiety.

- 4 -



RAPAMYCIN

It is preferred that the esters and ethers of rapamycin are of the hydroxyl groups at the 42- and/or 31-positions of the rapamycin nucleus, esters and ethers of a hydroxyl group at the 27-position (following chemical reduction of the 27-ketone), and that the oximes, hydrazones, and hydroxylamines are of a ketone at the 42-position (following oxidation of the 42-hydroxyl group) and of 27-ketone of the rapamycin nucleus.

10


Preferred 42- and/or 31-esters and ethers of rapamycin are disclosed in the following patents, which are all hereby incorporated by reference: alkyl esters (U.S. Patent 4,316,885); aminoalkyl esters (U.S. Patent 4,650,803); fluorinated esters (U.S. Patent 5,100,883); amide esters (U.S. Patent 5,118,677); carbamate esters (U.S. Patent 5,118,678); silyl ethers (U.S. Patent 5,120,842); aminoesters (U.S. Patent 5,130,307); acetals (U.S. Patent 5,51,413); aminodiester (U.S. Patent 5,162,333); sulfonate and sulfate esters (U.S. Patent 5,177,203); esters (U.S. Patent 5,221,670); alkoxyesters (U.S. Patent 5,233,036); O-aryl, -alkyl, -alkenyl, and -alkynyl ethers (U.S. Patent 5,258,389); carbonate esters (U.S. Patent 5,260,300); arylcarbonyl and alkoxy carbonyl carbamates (U.S. Patent 5,262,423); carbamates (U.S. Patent 5,302,584); hydroxyesters (U.S. Patent 5,362,718); hindered esters (U.S. Patent 5,385,908); heterocyclic esters (U.S. Patent 5,385,909); gem-disubstituted esters (U.S. Patent 5,385,910); amino alkanolic esters (U.S. Patent 5,389,639);

20

- 5 -

phosphorylcarbamate esters (U.S. Patent 5,391,730); carbamate esters (U.S. Patent 5,411,967); carbamate esters (U.S. Patent 5,434,260); amidino carbamate esters (U.S. Patent 5,463,048); carbamate esters (U.S. Patent 5,480,988); carbamate esters (U.S. Patent 5,480,989); carbamate esters (U.S. Patent 5,489,680); hindered N-oxide esters (U.S. Patent 5,491,231); biotin esters (U.S. Patent 5,504,091); O-alkyl ethers (U.S. Patent 5,665,772); and PEG esters of rapamycin (U.S. Patent 5,780,462). The preparation of these esters and ethers is disclosed in the patents listed above.

Accordingly examples of rapamycin compounds include compounds of
 10 formula:



wherein R^A and R^B are each selected from hydrogen and ester or ether forming groups as disclosed in any one of the abovementioned US patents.

15

Preferred 27-esters and ethers of rapamycin are disclosed in U.S. Patent 5,256,790, which is hereby incorporated by reference. The preparation of these esters and ethers is disclosed in the aforementioned patent.

20

Preferred oximes, hydrazones, and hydroxylamines of rapamycin are disclosed in U.S. Patents 5,373,014, 5,378,836, 5,023,264, and 5,563,145, which are hereby incorporated by reference. The preparation of these oximes, hydrazones, and

- 6 -

hydroxylamines are disclosed in the above listed patents. The preparation of 42-oxorapamycin is disclosed in 5,023,263, which is hereby incorporated by reference.

Particularly preferred rapamycins include rapamycin [U.S. Patent 3,929,992],
5 rapamycin 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid [U.S. Patent 5,362,718], and 42-O-(2-hydroxy)ethyl rapamycin [U.S. Patent 5,665,772].

When applicable, pharmaceutically acceptable salts can be formed from organic and inorganic acids, for example, acetic, propionic, lactic, citric, tartaric,
10 succinic, fumaric, maleic, malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic, camphorsulfonic, and similarly known acceptable aids when the rapamycin contains a suitable basic moiety. Salts may also be formed from organic and inorganic bases, such as alkali metal salts (for example, sodium,
15 lithium, or potassium) alkaline earth metal salts, ammonium salts, alkylammonium salts containing 1-6 carbon atoms or dialkylammonium salts containing 1-6 carbon atoms in each alkyl group, and trialkylammonium salts containing 1-6 carbon atoms in each alkyl group, when the rapamycin contains a suitable acidic moiety.

20 As used in accordance with this invention, the term "providing," with respect to providing a compound or substance covered by this invention, means either directly administering such a compound or substance, or administering a prodrug, derivative or analog which will form an effective amount of the compound or substance within the body.

25

The ability of the rapamycins of this invention to treat or inhibit cardiovascular disease or peripheral vascular disease was confirmed in a standard pharmacological test procedure using ApoE knockout (EKO) mice, which are a well accepted animal model of human atherosclerosis. In this test procedure, rapamycin
30 was used a representative example of a rapamycin of this invention. The procedure used, and results obtained are briefly summarized below.

- 7 -

Male EKO mice, 4-6 weeks of age, were housed in shoe-box cages and were allowed ad lib food and water. The animals were randomized by weight into 5 groups (N=12-15 mice per group) and were fed Purina Rodent Chow for the first week of the study. Also during this period as well as the remaining 12 weeks of the study, the animals were dosed every 2 days with 0, 1, 2, 4 or 8 mg/kg rapamycin s.c. using 2% Tween-80, 1% carboxymethyl cellulose as the vehicle and Control. The animal diet was switched to a casein-based Western Diet for week 2 to week 13 of the study. At the end of the study period, the animals were euthanized, plasma samples obtained, and the hearts perfused first with saline, then with 10% formalin. Total cholesterol and triglycerides were determined using enzymatic methods with commercially-available kits from Boehringer Mannheim and Wako Biochemicals, respectively, and the Boehringer Mannheim Hitachi 911 Analyzer (Boehringer Mannheim Diagnostic Laboratory Systems, Indianapolis, IN). Separation and quantification of plasma lipoproteins were performed using FPLC size fractionation. Briefly, 50-100 ml of serum was filtered and injected into two Superose 6 columns (Amersham Pharmacia Biotech, UK, Ltd) connected in series and eluted at a constant flow rate with 1 mM sodium EDTA and 0.15 M NaCl. Areas of each curve representing VLDL, LDL and HDL were integrated using Millennium software (Waters Technologies Corporation), and each lipoprotein fraction was quantified by multiplying the Total Cholesterol value by the relative percent area of each respective peak. The aortas were carefully isolated and remained in the formalin fixative for 48-72 hours before handling. Atherosclerotic lesions were identified by Oil Red O staining. The vessels were destained, and then imaged using a Nikon SMU800 microscope fitted with a Sony 3CCD video camera system in concert with IMAQ Configuration Utility (National Instrument) as the image capturing software. The lesions were quantified along the aortic arch using a custom threshold utility software package designed by Robert Coll (Coleman Technologies). Automated lesion assessment was performed on the vessels using the threshold function of the program, specifically on the region contained within the aortic arch from the proximal edge of the Right Common Carotid artery to the distal edge of the Left Subclavian artery. Aortic atherosclerosis data were expressed as percent lesion involvement strictly within this defined luminal area. Statistical significance between the Control and treated groups was determined using the Dunnett's Test at 1% significance level ($p < 0.01$).

- 8 -

The following table summarizes the results obtained in this standard pharmacological test procedure for atherosclerosis.

Table 1. The Effect of Rapamycin on Plasma Lipids and Aortic Atherosclerosis in Apo E Deficient mice

<u>Dosage</u>	<u>Triglycerides</u> (mg/dl)	<u>Total Cholesterol</u> (mg/dl)	<u>VLDL-C</u> (mg/dl)	<u>LDL-C</u> (mg/dl)	<u>HDL-C</u> (mg/dl)	<u>Aortic Atherosclerosis</u> (% lesion involvement)
Control	104±13	1186±47	807±48	371±13	7±3	39.5±2.6
1 mg/kg ⁺	132±16	1434±35	903±34	508±18*	23±6	21.6±3.1*
2 mg/kg	143±20	1311±80	763±64	517±18*	31±5*	14.9±3.1*
4 mg/kg	136±12	1281±58	749±58	494±10*	38±5*	16.4±2.8*
8 mg/kg	134±9	1167±75	644±58	475±21*	49±3*	12.03±2.3*

- 5 Data are mean ± S.E.
 +Dosage of rapamycin.
 *Significantly different from Control group (p<0.01).

The results in Table I show that treatment with rapamycin significantly (p < 0.01) increased levels of HDL-cholesterol and LDL-cholesterol, while not significantly affecting levels of triglycerides, total cholesterol, and VLDL-cholesterol compared with control EKO mice. Table I also shows a marked and dramatic decrease in the level of atherosclerosis in the rapamycin treated mice. While animals of the Control group demonstrated a mean lesion involvement in the aortic arch of 39.6%, atherosclerosis in animals treated with rapamycin was only 21.6% involvement at 1 mg/kg and decreased further to 14%, 16%, and 12 % at the 2, 4, and 8 mg/kg dosages, respectively. This represents a dramatic three-fold reduction in aortic atherosclerosis in a well accepted model of human atherosclerosis.

Based on the results obtained in the standard pharmacological test procedure described above, rapamycins are useful in the treatment or inhibition of cardiovascular disease and peripheral vascular disease. More particularly, the rapamycins of this invention are useful in treating or inhibiting coronary artery disease, cerebrovascular disease, arteriosclerosis, atherosclerosis, nonatheromatous arteriosclerosis, or vascular wall damage from cellular events leading toward immune

- 9 -

mediated vascular damage. The rapamycins of this invention are also useful inhibiting stroke or multiinfarct dementia.

In accordance with this invention, contemplated that a rapamycin may be used
5 as the sole active ingredient to provide the cardiovascular, cerebral, or peripheral
vascular benefits covered by this invention, or may be administered in combination
with other agents which provide beneficial cardiovascular, cerebral, or peripheral
vascular effects. Such agents are generally in the classes of compounds known as
ACE inhibitors, such as quinapril, perindopril, ramipril, captopril, trandolapril,
10 fosinopril, lisinopril, moexipril, and enalapril; angiotensin II receptor antagonists,
such as candesartan, irbesartan, losartan, valsartan, and telmisartan; fibric acid
derivatives, such as clofibrate, and gemfibrozil; HMG Co-A reductase inhibitors, such
as cerivastatin, fluvastatin, atorvastatin, lovastatin, pravastatin, simvastatin; beta
adrenergic blocking agents, such as sotalol, timolol, esmolol, carteolol, propranolol,
15 betaxolol, penbutolol, nadolol, acebutolol, atenolol, metoprolol, and bisoprolol;
calcium channel blockers, such as nifedipine, verapamil, nicardipine, diltiazem,
nimodipine, amlodipine, felodipine, nisoldipine, and bepridil; antioxidants;
anticoagulants such as, warfarin, dalteparin, heparin, enoxaparin, and danaparoid; and
agents useful in hormone replacement therapy containing estrogens, such as
20 conjugated estrogens, ethinyl estradiol, 17-beta-estradiol, estradiol, and estropipate.

Accordingly this invention also provides a product comprising a rapamycin and
one or more agents selected from an ACE inhibitor, an angiotensin II receptor
antagonists, a fibric acid derivative, a HMG Co-A reductase inhibitor, a beta
25 adrenergic blocking agent, a calcium channel blocker, an antioxidant; an
anticoagulants, and an agent useful in hormone replacement therapy containing an
estrogen for administration as a combined preparation for simultaneous, separate or
sequential use in treating or inhibiting cardiovascular, cerebral vascular, or peripheral
vascular disease in a mammal.

30

It is understood that the effective dosage of a rapamycin may vary depending
upon the particular compound utilized, the mode of administration, the condition, and
severity thereof, of the condition being treated, as well as the various physical factors

- 10 -

related to the individual being treated. As used in accordance with invention, satisfactory results may be obtained when the rapamycin is administered in a daily oral dosage of from about 5 μ g to 0.75 mg per kilogram of body weight. The projected daily dosages are expected to vary with route of administration.

5

When a rapamycin is used as part of a combination regimen, dosages of each of the components of the combination are administered during a desired treatment period. The components of the combination may administered at the same time; either as a unitary dosage form containing both components, or as separate dosage
10 units; the components of the combination can also be administered at different times during during a treatment period, or one may be administered as a pretreatment for the other.

Such doses may be administered in any manner useful in directing the active
15 compounds herein to the recipient's bloodstream, including orally, via implants, parenterally (including intravenous, intraperitoneal and subcutaneous injections), rectally, intranasally, vaginally, and transdermally. For the purposes of this disclosure, transdermal administrations are understood to include all administrations across the surface of the body and the inner linings of bodily passages including
20 epithelial and mucosal tissues. Such administrations may be carried out using the present compounds, or pharmaceutically acceptable salts thereof, in lotions, creams, foams, patches, suspensions, solutions, and suppositories (rectal and vaginal).

Oral formulations containing the active compounds of this invention may
25 comprise any conventionally used oral forms, including tablets, capsules, buccal forms, troches, lozenges and oral liquids, suspensions or solutions. Capsules may contain mixtures of the active compound(s) with inert fillers and/or diluents such as the pharmaceutically acceptable starches (e.g. corn, potato or tapioca starch), sugars, artificial sweetening agents, powdered celluloses, such as crystalline and
30 microcrystalline celluloses, flours, gelatins, gums, etc. Useful tablet formulations may be made by conventional compression, wet granulation or dry granulation methods and utilize pharmaceutically acceptable diluents, binding agents, lubricants, disintegrants, surface modifying agents (including surfactants), suspending or

- 11 -

stabilizing agents, including, but not limited to, magnesium stearate, stearic acid, talc, sodium lauryl sulfate, microcrystalline cellulose, carboxymethylcellulose calcium, polyvinylpyrrolidone, gelatin, alginic acid, acacia gum, xanthan gum, sodium citrate, complex silicates, calcium carbonate, glycine, dextrin, sucrose, sorbitol, dicalcium phosphate, calcium sulfate, lactose, kaolin, mannitol, sodium chloride, talc, dry starches and powdered sugar. Preferred surface modifying agents include nonionic and anionic surface modifying agents. Representative examples of surface modifying agents include, but are not limited to, poloxamer 188, benzalkonium chloride, calcium stearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, magnesium aluminum silicate, and triethanolamine. It is more preferred that poloxamer 188 is used as the surface modifying agent. Oral formulations herein may utilize standard delay or time release formulations to alter the absorption of the active compound(s). Preferred oral formulations of rapamycins are disclosed in U.S. Patents 5,559,121; 5,536,729; 5,989,591; and 5,985,325, which are hereby incorporated by reference.

In some cases it may be desirable to administer the compounds directly to the airways in the form of an aerosol.

The compounds of this invention may also be administered parenterally or intraperitoneally. Solutions or suspensions of these active compounds as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxy-propylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can

- 12 -

be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils. Preferred parenteral formulations for administering a rapamycin are disclosed in U.S. Patents 5,530,006; 5,516,770; and 5,616,588, which
5 are hereby incorporated by reference.

Suppository formulations may be made from traditional materials, including cocoa butter, with or without the addition of waxes to alter the suppository's melting point, and glycerin. Water soluble suppository bases, such as polyethylene glycols of
10 various molecular weights, may also be used.

- 13 -

CLAIMS

1. A method of treating or inhibiting cardiovascular, cerebral vascular, or peripheral vascular disease in a mammal in need thereof, which comprises providing
5 said mammal with an effective amount of a rapamycin.
2. The method according to claim 1, wherein the rapamycin is rapamycin.
3. The method according to claim 1, wherein the rapamycin is a ester,
10 ether, oxime, hydrazone, or hydroxylamine of rapamycin
4. The method according to claim 3, wherein the rapamycin is a 42-ester or 42-ether of rapamycin.
- 15 5. The method according to claim 4, wherein the rapamycin is rapamycin 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid.
6. The method according to claim 4, wherein the rapamycin is 42-O-(2-hydroxy)ethyl rapamycin.
20
7. The method according to any one of claims 1 to 6 wherein the rapamycin is provided in combination with one or more agents selected from the groups consisting of an ACE inhibitor, an angiotensin II receptor antagonists, a fibric acid derivative, a HMG Co-A reductase inhibitor, a beta adrenergic blocking agent, a
25 calcium channel blocker, an antioxidant; an anticoagulants, or an agent useful in hormone replacement therapy containing an estrogen.
8. A method of treating or inhibiting coronary artery disease, cerebrovascular disease, arteriosclerosis, atherosclerosis, nonatheromatous arterio-
30 sclerosis, or vascular wall damage from cellular events leading toward immune mediated vascular damage in a mammal in need thereof, which comprises providing said mammal with an effective amount of a rapamycin.

- 14 -

9. The method according to claim 8, wherein the rapamycin is rapamycin.

10. The method according to claim 8, wherein the rapamycin is a ester, ether, oxime, hydrazone, or hydroxylamine of rapamycin

5

11. The method according to claim 10, wherein the rapamycin is a 42-ester or 42-ether of rapamycin.

12. The method according to claim 11, wherein the rapamycin is rapamycin
10 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid.

13. The method according to claim 11, wherein the rapamycin is 42-O-(2-hydroxy)ethyl rapamycin.

14. The method according to any one of claims 8 to 13 wherein the
15 rapamycin is provided in combination with one or more agents selected from the groups consisting of an ACE inhibitor, an angiotensin II receptor antagonists, a fibric acid derivative, a HMG Co-A reductase inhibitor, a beta adrenergic blocking agent, a calcium channel blocker, an antioxidant; an anticoagulants, or an agent useful in
20 hormone replacement therapy containing an estrogen.

15. A method of inhibiting stroke or multiinfarct dementia in a mammal in need thereof, which comprises providing said mammal with an effective amount of a rapamycin.

25

16. The method according to claim 15, wherein the rapamycin is rapamycin.

17. The method according to claim 15, wherein the rapamycin is a ester,
30 ether, oxime, hydrazone, or hydroxylamine of rapamycin

18. The method according to claim 17, wherein the rapamycin is a 42-ester or 42-ether of rapamycin.

- 15 -

19. The method according to claim 18, wherein the rapamycin is rapamycin 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid.

20. The method according to claim 18, wherein the rapamycin is 42-O-(2-hydroxy)ethyl rapamycin.

21. The method according to any one of claims 15 to 20 wherein the rapamycin is provided in combination with one or more agents selected from the groups consisting of an ACE inhibitor, an angiotensin II receptor antagonists, a fibric acid derivative, a HMG Co-A reductase inhibitor, a beta adrenergic blocking agent, a calcium channel blocker, an antioxidant; an anticoagulants, or an agent useful in hormone replacement therapy containing an estrogen.

22. Use of a rapamycin in the preparation of a medicament for treating or inhibiting cardiovascular, cerebral vascular, or peripheral vascular disease in a mammal.

23. Use of a rapamycin in the preparation of a medicament for treating or inhibiting coronary artery disease, cerebrovascular disease, arteriosclerosis, atherosclerosis, nonatheromatous arteriosclerosis, or vascular wall damage from cellular events leading toward immune mediated vascular damage in a mammal.

24. Use of a rapamycin in the preparation of a medicament for inhibiting stroke or multiinfarct dementia in a mammal.

25. A product comprising a rapamycin and one or more agents selected from an ACE inhibitor, an angiotensin II receptor antagonists, a fibric acid derivative, a HMG Co-A reductase inhibitor, a beta adrenergic blocking agent, a calcium channel blocker, an antioxidant; an anticoagulants, and an agent useful in hormone replacement therapy containing an estrogen for administration as a combined preparation for simultaneous, separate or sequential use in treating or inhibiting cardiovascular, cerebral vascular, or peripheral vascular disease in a mammal.

- 16 -

26. A product comprising a rapamycin and one or more agents selected from an ACE inhibitor, an angiotensin II receptor antagonists, a fibric acid derivative, a HMG Co-A reductase inhibitor, a beta adrenergic blocking agent, a calcium channel blocker, an antioxidant; an anticoagulants, and an agent useful in hormone replacement therapy containing an estrogen for administration as a combined preparation for simultaneous, separate or sequential use in treating or inhibiting coronary artery disease, cerebrovascular disease, arteriosclerosis, atherosclerosis, nonatheromatous arteriosclerosis, or vascular wall damage from cellular events leading toward immune mediated vascular damage in a mammal.

10

27. A product comprising a rapamycin and one or more agents selected from an ACE inhibitor, an angiotensin II receptor antagonists, a fibric acid derivative, a HMG Co-A reductase inhibitor, a beta adrenergic blocking agent, a calcium channel blocker, an antioxidant; an anticoagulants, and an agent useful in hormone replacement therapy containing an estrogen for administration as a combined preparation for simultaneous, separate or sequential use in inhibiting stroke or multiinfarct dementia in a mammal.

15

28. A product according to any one of claims 25 to 27 wherein the rapamycin is rapamycin.

20

29. A product according to any one of claims 25 to 27 wherein the rapamycin is a ester, ether, oxime, hydrazone, or hydroxylamine of rapamycin

25

30. A product according to any one of claims 25 to 27 wherein the rapamycin is a 42-ester or 42-ether of rapamycin.

30

31. A product according to any one of claims 25 to 27 wherein the rapamycin is rapamycin 42-ester with 3-hydroxy-2-(hydroxymethyl)-2-methylpropionic acid.

32. A product according to any one of claims 25 to 27 wherein the rapamycin is 42-O-(2-hydroxy)ethyl rapamycin.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2001 (27.12.2001)

PCT

(10) International Publication Number
WO 01/97809 A3

- (51) International Patent Classification⁷: **A61K 31/436**, 45/06, A61P 9/00, 9/10, 25/28, A61K 31/435 // (A61K 45/06, 31:71, 31:445)
- (21) International Application Number: **PCT/US01/19179**
- (22) International Filing Date: **14 June 2001 (14.06.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/212.117 **16 June 2000 (16.06.2000)** **US**
- (71) Applicant: **AMERICAN HOME PRODUCTS CORPORATION** [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US).
- (72) Inventors: **AZROLAN, Neal, Ivan**; 286 Glenn Avenue, Lawrenceville, NJ 08648 (US). **SEHGAL, Surendra, Nath**; 20711 76th Avenue S.E., Snohomish, WA 98296 (US). **ADELMAN, Steven, Jay**; 342 Windy Run Drive, Doylestown, PA 18901 (US).
- (74) Agents: **MILOWSKY, Arnold, S.**; American Home Products Corporation, Patent Law Dept., Five Giralda Farms, Madison, NJ 07940 et al. (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW). Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR). OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*
- (88) Date of publication of the international search report:
10 May 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/97809 A3

(54) Title: **METHOD OF TREATING CARDIOVASCULAR DISEASE USING RAPAMYCIN**

(57) Abstract: The invention provides a method of treating or inhibiting cardiovascular, cerebral vascular, or peripheral vascular disease in a mammal in need thereof, which comprises providing said mammal an effective amount of a rapamycin.

INTERNATIONAL SEARCH REPORT

Inte: onal Application No
PCT/US 01/19179

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/436 A61K45/06 A61P9/00 A61P9/10 A61P25/28
A61K31/435 //(A61K45/06, 31:71, 31:445)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS, EMBASE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 015 809 A (T. ZHU, H-K. LEE) 18 January 2000 (2000-01-18) claim 1 column 1, line 13-20 column 5, line 1-7 ---	1-4, 8-11, 22, 23
X	US 5 373 014 A (A.A. FAILLI E.A.) 13 December 1994 (1994-12-13) cited in the application claims 1,4 column 1, line 5-11 column 6, line 10-17 --- -/--	1-4, 8-11, 22, 23



Further documents are listed in the continuation of box C



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

8 document member of the same patent family

Date of the actual completion of the international search

26 February 2002

Date of mailing of the international search report

04/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Peeters, J

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/19179

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 516 781 A (R.E. MORRIS, C.R. GREGORY) 14 May 1996 (1996-05-14) cited in the application claim 1 column 10, line 34-39 ---	1,2,8,9, 22,23
X	WO 97 35575 A (NOVARTIS) 2 October 1997 (1997-10-02) claims 1,3,5-7 ---	1-3,8,9, 22,23
X	WO 98 56358 A (AMERICAN HOME PRODUCTS) 17 December 1998 (1998-12-17) claim 1 page 1, line 3-7 ---	1,2,8,9, 22,23
X	WO 96 17845 A (AMERICAN HOME PRODUCTS) 13 June 1996 (1996-06-13) claims 1,19 -----	1-3,8,9, 22,23

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/19179

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6015809	A	18-01-2000	US 6117863 A	12-09-2000
US 5373014	A	13-12-1994	US 5446048 A	29-08-1995
US 5516781	A	14-05-1996	AT 210979 T	15-01-2002
			AU 706486 B2	17-06-1999
			AU 1791895 A	23-11-1995
			CA 2148871 A1	13-11-1995
			DE 69524679 D1	31-01-2002
			EP 0691130 A1	10-01-1996
			JP 8040901 A	13-02-1996
			NZ 272092 A	27-07-1997
			US 5665728 A	09-09-1997
			US 5646160 A	08-07-1997
			US 5563146 A	08-10-1996
			AT 194493 T	15-07-2000
			AU 665041 B2	14-12-1995
			AU 3107993 A	15-07-1993
			CA 2086642 A1	10-07-1993
			DE 69328981 D1	17-08-2000
			DE 69328981 T2	16-11-2000
			DK 551182 T3	09-10-2000
			EP 0551182 A1	14-07-1993
			ES 2147740 T3	01-10-2000
			GR 3034479 T3	29-12-2000
			HK 1010332 A1	29-12-2000
			JP 2644955 B2	25-08-1997
			JP 6009390 A	18-01-1994
			LV 12603 A	20-01-2001
			LV 12603 B	20-04-2001
			NZ 245630 A	27-07-1997
			PT 551182 T	29-12-2000
			SG 47639 A1	17-04-1998
WO 9735575	A	02-10-1997	AU 716514 B2	24-02-2000
			AU 2291897 A	17-10-1997
			BR 9708358 A	03-03-1999
			CA 2247275 A1	02-10-1997
			CN 1214631 A	21-04-1999
			CZ 9803070 A3	13-01-1999
			WO 9735575 A1	02-10-1997
			EP 1149581 A2	31-10-2001
			EP 0893996 A1	03-02-1999
			HU 0001267 A2	28-11-2000
			JP 2000507226 T	13-06-2000
			NO 984340 A	18-09-1998
			PL 328659 A1	15-02-1999
			SK 132798 A3	10-03-1999
			TR 9801919 T2	22-02-1999
			ZA 9702719 A	28-09-1998
WO 9856358	A	17-12-1998	AU 7836898 A	30-12-1998
			BR 9810746 A	19-09-2000
			CN 1259864 T	12-07-2000
			EP 0994697 A1	26-04-2000
			HU 0003341 A2	28-02-2001
			NO 996044 A	08-12-1999
			PL 337320 A1	14-08-2000

INTERNATIONAL SEARCH REPORT

...formation on patent family members

International Application No

PCT/US 01/19179

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9856358	A		SK 171799 A3	11-07-2000
			TR 9903065 T2	21-09-2000
			WO 9856358 A1	17-12-1998
			ZA 9805095 A	13-12-1999
<hr/>				
WO 9617845	A	13-06-1996	US 5563145 A	08-10-1996
			AT 202357 T	15-07-2001
			AU 714120 B2	16-12-1999
			AU 4417696 A	26-06-1996
			BR 9509875 A	30-09-1997
			CA 2206965 A1	13-06-1996
			CN 1174554 A	25-02-1998
			CZ 9701720 A3	14-01-1998
			DE 69521438 D1	26-07-2001
			DK 796260 T3	03-09-2001
			EP 0796260 A1	24-09-1997
			ES 2157353 T3	16-08-2001
			HU 77141 A2	02-03-1998
			JP 10510270 T	06-10-1998
			PT 796260 T	28-09-2001
			WO 9617845 A1	13-06-1996
			US 5677295 A	14-10-1997